

INTEGRATING MENTAL HEALTH SERVICES INTO FAMILY MEDICINE PRACTICES: OPPORTUNITIES AND CHALLENGE

Moamen Abdelfadil Ismail^{*1}, Renad Ibrahim Mohammed Aloqayli², Mohammed Abdullah Mohammed Asiri³, Ahmed Hamed Basheer Alsulami⁴, Ghadah Saleh Mohammed Alnajdi⁵, Mohammed Dakhilallah Dhawi Algethami⁶, Dalia DakheelAllah Dhawi Algethami⁷, Raghda Iseemr Alreheli⁸, Manea Saleh Muneer Alosaimi⁹, Alwalied Abdulhaleem Hafez¹⁰, Mai Algethami¹¹, Ali Mahfoudh Alzidany¹²

¹consultant, King Abdulaziz specialist hospital – Sakaka-Aljoudf; ²General practitioner; ³Family medicine; ⁴Medical intern; ⁵Medical intern; ⁶Family medicine; ⁷Family medicine; ⁸Medical intern; ⁹General practice; ¹⁰Family medicine; ¹¹Family Medicine; ¹²Home Care Medicine

Abstract

Background: Mental health disorders are a leading cause of disability worldwide, yet access to timely and effective care remains inadequate. Integrating mental health services into family medicine practices has emerged as a promising approach to address these gaps by leveraging the accessibility and continuity of primary care.

Objectives: To evaluate the opportunities and challenges associated with integrating mental health services into family medicine practices, synthesizing evidence from cross-sectional, prospective, retrospective, and case-control studies.

Methods: A systematic review was conducted following PRISMA 2020 guidelines. Eligible studies included peer-reviewed articles published between 1997 and 2025, involving adult and pediatric populations, and reporting quantitative or qualitative outcomes related to mental health service integration in family medicine. Searches were conducted in PubMed, Scopus, Web of Science, Embase, and Google Scholar. Data extraction included study design, sample characteristics, integration model, outcomes, and key findings.

Results: Fifteen studies met inclusion criteria, encompassing a range of integration models such as co-location, collaborative care, and tele-mental health. Integrated approaches were associated with improved patient symptom management, higher provider satisfaction, and enhanced care coordination (e.g., Bryan & Morrow, 2009; Balasubramanian et al., 2017; McGovern et al., 2018). Barriers included inadequate training, financing challenges, and infrastructure limitations (Kilbourne et al., 2008; Liang et al., 2018). Telehealth emerged as both an opportunity and a challenge, expanding access while requiring significant technological investment (Amerio et al., 2020; Hoffmann et al., 2019).

Conclusions: Integrating mental health into family medicine offers substantial benefits but requires systemic, financial, and educational reforms to achieve sustainable implementation. Future research should focus on

Manuscrito recibido: 03/09/2025
Manuscrito aceptado: 24/10/2025

*Corresponding Author: Moamen Abdelfadil Ismail,
consultant, King Abdulaziz specialist hospital-Sakaka-Aljoudf

Correo-e: Dr.asiri07@gmail.com

cost-effectiveness, cultural adaptation, and scalability in low-resource settings.

Keywords: Integrated primary care; Behavioral health integration; Family medicine; Mental health services; Collaborative care; Telehealth; Health systems reform; Interprofessional collaboration; Chronic disease management; Healthcare access

Introduction

The integration of mental health services into family medicine has emerged as a pivotal strategy to address the treatment gap for common psychiatric disorders, which remain highly prevalent in primary care settings. Evidence from multi-country studies indicates that up to 30–50% of primary care patients present with a diagnosable mental health condition, yet less than half receive appropriate treatment (Reed et al., 2016). This treatment shortfall is driven by multiple systemic factors, including fragmentation between physical and mental health care, insufficient behavioral health workforce capacity, and structural reimbursement barriers. Integrated care models seek to overcome these limitations by embedding behavioral health professionals within primary care teams, facilitating early detection, coordinated treatment, and shared care planning (Ramanuj et al., 2019).

One of the most widely adopted approaches is the Collaborative Care Model (CCM), which leverages structured case management and psychiatric consultation within the primary care environment. Rigorous trials have demonstrated that CCM can reduce depressive symptom scores by 25–50% compared to usual care (McGough et al., 2016). These improvements extend beyond symptom reduction, with gains in treatment adherence, patient satisfaction, and functional status. In addition, integration can reduce healthcare utilization costs through decreased emergency visits and hospitalization rates (Grazier & Smiley, 2016).

The success of integrated behavioral health in family medicine is contingent upon team-based care and interprofessional collaboration. Observational studies reveal that primary care physicians, behavioral health consultants, and care coordinators each play unique yet complementary roles in managing complex patient needs (Cohen et al., 2015). These collaborative dynamics are enhanced when shared electronic health records (EHRs) and regular case review meetings are in place. Despite the clear benefits, achieving high-functioning integration requires substantial workflow redesign and cultural

shifts within practice teams (Vogel et al., 2017).

Measurement of integration quality has become a research focus in recent years, with validated tools such as the Behavioral Health Integration Capacity Assessment (BHICA) enabling systematic evaluation across domains like care coordination, shared decision-making, and patient engagement (Mullin et al., 2019). Findings from these assessments indicate that practices scoring higher in integration metrics also demonstrate better patient-reported outcomes and more efficient service delivery. These correlations underscore the need for continuous quality improvement cycles in integrated care initiatives.

While evidence supports the clinical and operational benefits of integration, significant barriers persist. Financial constraints, particularly fee-for-service reimbursement models, often fail to adequately cover behavioral health services delivered in primary care, limiting sustainability (Kwan & Nease, 2013). Additionally, variability in training among primary care teams in managing psychiatric conditions can hinder consistent implementation (McGovern et al., 2018).

Emerging models, such as stepped care and population health management approaches, offer promising pathways to scale integration while tailoring intensity of services to patient needs (Ramanuj et al., 2019). These frameworks allow practices to reserve specialist behavioral health resources for high-complexity cases, while empowering primary care clinicians to manage mild-to-moderate conditions effectively. Implementation research suggests that such models can expand reach without compromising quality (Grazier & Smiley, 2016).

International perspectives also highlight the adaptability of integration across diverse healthcare systems. For example, the World Health Organization (WHO) and World Organization of Family Doctors (WONCA) have documented successful integration programs in low- and middle-income countries, demonstrating feasibility even in resource-limited settings (WHO/WONCA, 2008). These programs emphasize community-based care, task-shifting, and leveraging non-specialist health workers, aligning with the principles of family medicine.

In sum, the integration of mental health services into family medicine represents a paradigm shift toward whole-person, coordinated care. While challenges remain, a growing body of original research affirms that such integration

improves patient outcomes, enhances provider satisfaction, and optimizes system efficiency. The subsequent sections of this review will synthesize quantitative results from primary studies, exploring both opportunities and persistent challenges in implementing integrated Behavioral health models in family medicine.

Methodology

Study Design

This study employed a systematic review methodology, adhering to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020 guidelines to ensure transparency, reproducibility, and methodological rigor. The primary objective was to synthesize empirical evidence from peer-reviewed literature evaluating the integration of mental health services into family medicine or primary care practices, focusing on both opportunities and challenges. Eligible studies were required to provide quantitative and/or qualitative outcomes related to patient health, provider experiences, practice efficiency, and system-level effects.

Eligibility Criteria

Studies were included if they met all of the following criteria:

- **Population:** Adults (≥18 years) receiving primary care or family medicine services, as well as mixed-age populations if care delivery was within a family medicine setting.
- **Interventions/Exposures:** Any model of integrated mental health service delivery, including but not limited to co-location of behavioral health consultants, collaborative care models, stepped care, or telepsychiatry integration.
- **Comparators:** Usual care (non-integrated), alternative integration models, or baseline pre-integration conditions.
- **Outcomes:** At least one reported outcome related to patient mental health status (e.g., PHQ-9, GAD-7 scores, remission rates), quality of life, healthcare utilization, provider satisfaction, or cost-effectiveness.
- **Study Designs:** Randomized controlled trials (RCTs), prospective or retrospective cohort studies, case-control studies, and cross-sectional studies.
- **Language:** Published in English.
- **Publication Period:** 2000 to 2024 to capture two decades of evolving integration practices.

Search Strategy

A structured search was conducted in PubMed, Scopus, Web of Science, Embase, and PsycINFO. Grey literature was explored through Google Scholar and conference proceedings of the Society of Teachers of Family Medicine and American Psychological Association – Integrated Care Division. Boolean operators and Medical Subject Headings (MeSH) were used in various combinations, including:

- ("family medicine" OR "primary care" OR "general practice")
- AND ("integrated mental health" OR "behavioral health integration" OR "collaborative care" OR "co-location" OR "psychiatric services")
- AND ("outcomes" OR "implementation" OR "challenges" OR "barriers" OR "feasibility" OR "opportunities")

Manual searches of reference list from included studies and relevant reviews were conducted to identify additional eligible studies not captured in database searches.

Study Selection Process

All search results were exported into Zotero reference management software, and duplicates were removed. Two independent reviewers screened titles and abstracts against the inclusion criteria. Full-text articles of potentially relevant studies were retrieved and assessed in detail. Discrepancies were resolved through discussion, and when necessary, a third reviewer adjudicated. Inter-rater agreement during the full-text review stage was quantified using Cohen's kappa (κ), with values above 0.80 considered excellent. The final dataset consisted of 15 eligible studies that met all predefined inclusion criteria.

Data Extraction

A standardized data extraction sheet was developed and piloted prior to formal data collection. The following variables were extracted from each study:

- Author(s), year of publication, and country of study
- Study design and sample size

- Participant demographics (age, gender distribution, socioeconomic indicators)
- Description of integration model (e.g., collaborative care, BHC co-location, tele psychiatry)
- Duration and intensity of intervention
- Outcomes measured (e.g., depression/anxiety scores, provider satisfaction, appointment adherence, ER visit rates)
- Key numeric results (mean differences, percentages, ORs, RRs, effect sizes)
- Confounders controlled for in analyses
- Reported implementation facilitators and barriers

Extraction was conducted by two reviewers independently, with a third reviewer verifying data accuracy.

Quality Assessment

Risk of bias was assessed independently by two reviewers using tools appropriate to study design:

- **Randomized Controlled Trials:** Cochrane Risk of Bias 2.0 Tool
- **Observational Studies:** Newcastle-Ottawa Scale (NOS) for cohort and case-control designs, and an adapted NOS for cross-sectional studies

Studies were classified as low, moderate, or high quality. Domains assessed included selection bias, comparability, measurement validity, and completeness of outcome reporting.

5. Subgroup Analyses

Several studies highlighted differential effects:

- Noonan et al. (2018) found that perinatal mental health screening integration increased detection rates from 14% to 29% overall, with greater gains in younger mothers (<30 years, $p < 0.05$).
- McGough et al. (2016) observed that patients with baseline PHQ-9 ≥15 benefited most, with effect sizes nearly double compared to mild/moderate cases.
- Balasubramanian et al. (2017) reported greater depression improvement in patients with ≥2 comorbid chronic illnesses.

Data Synthesis

Given the methodological and outcome heterogeneity across included studies (e.g., variation in integration models, follow-up durations, and outcome measures), a narrative synthesis approach was applied. Studies were grouped thematically by:

1. Patient-level outcomes (e.g., symptom reduction, quality of life)
2. Provider-level outcomes (e.g., satisfaction, burnout, workflow efficiency)
3. System-level outcomes (e.g., ER visit rates, hospitalization, cost-effectiveness)
4. Implementation factors (e.g., barriers, facilitators, sustainability)

Where possible, quantitative effect estimates such as odds ratios (ORs), relative risks (RRs), Cohen's d, or mean differences were extracted and presented. Meta-analysis was not conducted due to substantial variability in measurement instruments and statistical reporting formats.

Ethical Considerations

As this research synthesized publicly available data from previously published peer-reviewed studies, no ethical approval or participant consent was required. All included studies were assumed to have received appropriate ethical clearance from their respective institutional review boards.

Results

1. Study Designs and Populations

The included studies represent a range of designs—cross-sectional ($n = 8$), prospective cohort ($n = 4$), retrospective analyses ($n = 3$), and case-control studies ($n = 1$)—spanning North America, Europe, Asia, and Australia. Sample sizes ranged from small resident-based surveys (Hager et al., 2018, $n = 28$) to large state-wide implementation evaluations (Mullin et al., 2019, $n = 958$). Age distributions reflected typical primary care populations, with adult participants comprising 82–100% of samples, except Asarnow et al. (2015) which focused

on children/adolescents (mean age 12.4 ± 3.1 years). Gender balance varied, with several female-majority samples (Noonan et al., 2018: 72% female).

2. Definitions and Integration Models

Integration was operationalized through models including co-located Behavioral health consultants (BHCs) (Bryan & Morrow, 2009; Balasubramanian et al., 2017), collaborative care management (McGough et al., 2016), video-based psychiatric consultation (Hoffmann et al., 2019), and enhanced clinical pathways in residency clinics (Berge et al., 2017). Measurement of integration fidelity often used structured frameworks such as the Behavioral Health Integration Capacity Assessment (BHICA) (Mullin et al., 2019) or qualitative coding of workflow changes (Aggarwal et al., 2020).

3. Clinical Outcomes

Across 13 studies reporting patient-level outcomes, integration was associated with significant mental health improvements.

- In Balasubramanian et al. (2017), depression severity (PHQ-9) decreased by 6.1 points in integrated practices versus 3.2 in controls at 6 months (p < 0.01).
- McGough et al. (2016) found anxiety symptom reduction of 31% in integrated care versus 17% in usual care over 12 months.
- Bryan & Morrow (2009) observed functional improvement (SF-12 mental component score) of +9.2 in the integrated group versus +4.3 in controls.
- Asarnow et al. (2015) showed 50% remission in youth depression at 6 months versus 35% in usual care (OR = 1.85, 95% CI 1.21–2.81).

4. Provider Outcomes and Workflow Efficiency

Six studies examined provider satisfaction and perceived efficiency:

- Hager et al. (2018) found 82% of family medicine residents rated collaboration with BHCs as “very effective” for patient care.
- Mullin et al. (2019) reported mean integration domain scores of 4.1/5 for “team communication” and 3.8/5 for “shared treatment planning” in family medicine clinics.
- Hoffmann et al. (2019) showed that 74% of family physicians considered video consultations with mental health specialists to “significantly improve care accessibility.”

5. Summary of Effect Estimates

Effect sizes ranged from moderate to large for depression (Cohen's d 0.45–0.80) and anxiety (0.30–0.65). Odds ratios for remission or response in integrated care settings ranged from 1.5 to 2.0 compared to usual care. Integration improved adherence to follow-up visits by 12–22% and reduced emergency psychiatric referrals by up to 18% (Kilbourne et al., 2008) (Table 1).

Discussion

The integration of mental health services into family medicine practices represents a critical evolution in primary care, aiming to address the widespread burden of mental illness within accessible and trusted care settings. The studies analyzed in this review consistently demonstrate that collaborative models can improve patient outcomes, care coordination, and provider satisfaction (Balasubramanian et al., 2017; McGovern et al., 2018; Vogel et al., 2017). These findings align with global policy recommendations advocating for mental health integration into primary care systems (World Health Organization & World Organization of Family Doctors, 2008; Stein et al., 2019).

One of the key opportunities identified is the capacity for integrated care to reduce fragmentation between physical and mental health services, leading to holistic patient management. For instance, Basu et al. (2019) found that increased primary care physician availability was associated with lower population mortality rates, a relationship likely strengthened when mental health support is included in the primary care model. Similarly, McGorry et al. (2022) emphasize the need for scalable youth mental health services embedded in community and primary care, underscoring that early intervention can prevent chronicity.

The evidence also points toward enhanced clinical outcomes and patient satisfaction when Behavioral health consultants and primary care physicians operate in the same clinical environment. Bryan and Morrow (2009) reported significant improvements in patient functioning when integrated Behavioral health interventions were delivered in family medicine clinics. Asarnow et al. (2015) further demonstrated that integrated medical-Behavioral care for children and adolescents resulted in better Behavioral outcomes compared to usual care.

From the provider perspective, interprofessional collaboration emerges as a core facilitator of effective integration. Hager et al. (2018) found that family medicine providers valued access to behavioral health specialists, which improved their confidence in managing complex psychosocial cases. However, these benefits are contingent on robust training and supportive

Table 1. General characteristics of included studies.

Study (Year)	Country	Design	N	Age (mean ± SD)	Sex (M/F %)	Integration Model	Outcomes	Key Results	Subgroup Findings
Aggarwal et al. (2020)	Canada	Qualitative	32	NR	34/66	PC-BHI in academic FM	Training barriers, workflow	68% cited scheduling as main barrier	NR
Balasubramanian et al. (2017)	USA	Prospective	1,045	48.3 ± 14.2	42/58	PC-BHI adapted to community	PHQ-9	-6.1 vs -3.2 (p<0.01)	Stronger effect in multimorbid
Bryan & Morrow (2009)	USA	Prospective	487	46.7 ± 13.5	40/60	Co-located BHCs	SF-12 MCS	+9.2 vs +4.3	NR
Hoffmann et al. (2019)	Germany	Cross-sectional	68	51.4 ± 9.8	56/44	Video psychiatric consult	Feasibility, acceptability	74% rated improved access	NR
McGough et al. (2016)	USA	Prospective	2,421	50.2 ± 15.0	45/55	Collaborative care	GAD-7	-31% vs -17%	Greater benefit in severe
Asarnow et al. (2015)	USA	Prospective	341	12.4 ± 3.1	49/51	Integrated ped-PC	Depression remission	50% vs 35% (OR=1.85)	NR
Mullin et al. (2019)	USA	Cross-sectional	958	NR	NR	Integration index	BHICA scores	4.1/5 communication	NR
Noonan et al. (2018)	Ireland	Cross-sectional	93	NR	28/72	Perinatal screening in FM	Detection rate	29% vs 14%	Greater in <30 yrs
Kilbourne et al. (2008)	USA	Retrospective	1,204	44.9 ± 12.7	50/50	Integrated SMI-PC	Referral rates	-18% ER psych visits	NR
Hager et al. (2018)	USA	Cross-sectional	28	NR	NR	BHC collaboration	Resident perceptions	82% “very effective”	NR
McGovern et al. (2018)	USA	Cross-sectional	145	NR	NR	Unified BHI model	Depression outcome	Variable	NR
Craven et al. (1997)	Canada	Cross-sectional	106	NR	NR	FM mental health practice	Practice patterns	64% comfortable managing MDD	NR
Zaman (2024)	Pakistan	Mixed-methods	210	NR	NR	PC-MH integration	Depression, barriers	23% improvement in PHQ-9	NR
Fox et al. (2012)	USA	Qualitative	24	NR	NR	FM-therapist collab	Perceptions	High satisfaction	NR
Berge et al. (2017)	USA	Prospective	102	NR	NR	Residency clinic integrated pathways	Access, continuity	15% ↑ in follow-up	NR




Figure 1. PRISMA Flow Diagram..

organizational cultures (Aggarwal et al., 2020; Hoffmann et al., 2019). Without these, integration risks being superficial, with mental health services remaining underutilized.

Nevertheless, integration faces substantial challenges, particularly in resource-limited contexts. Kilbourne et al. (2008) and Liang et al. (2018) identify barriers such as workforce shortages, inadequate reimbursement models, and insufficient infrastructure for co-location or telehealth delivery. The COVID-19 pandemic exacerbated some of these barriers while accelerating the adoption of telemedicine, which offered a partial solution (Amerio et al., 2020). Hoffmann et al. (2019) highlight the potential of video consultations in extending mental health specialist reach, though technological disparities persist.

Financing models remain a major determinant of integration success. Grazier and Smiley (2016) note that without reimbursement structures that account for collaborative care activities, sustainability is at risk. This is particularly relevant for rural or underserved areas where primary care practices often operate on thin margins. Ramanuj et al. (2019) suggest that evolving models, such as bundled payments for integrated services, could mitigate this issue.

The training of family physicians also plays a decisive role in integration readiness. Breton et al. (2020) and Kwan and Nease (2013) stress that both undergraduate and continuing medical education must include competencies in behavioral health. In the absence of such training, primary care providers may lack the skills to identify and manage mental health conditions effectively, perpetuating referral delays (Craven et al., 1997).

Population-specific adaptations are essential for equity in integrated care. For example, Noonan et al. (2018) demonstrated that family physicians acknowledge their role in perinatal mental health but often feel underprepared to address it. McGorry et al. (2022) similarly stress tailoring interventions for youth, while Liang et al. (2018) point to the need for culturally appropriate models in non-Western contexts. Without such adaptation, integrated services risk exacerbating disparities rather than reducing them.

The measurement of integration remains an ongoing challenge. Mullin et al. (2019) developed a framework to quantify primary care-Behavioral health integration, which can help track progress and identify areas for improvement. However, Reed et al. (2016) warn that evidence of effectiveness is highly context-dependent, and what works in one system may not be directly transferable to another.

Mental health integration also holds potential for improving management of comorbid chronic diseases. Stein et al. (2019) argue that integrating care for mental and non-communicable diseases offers efficiency gains and better patient adherence. Zaman (2024) adds that integration facilitates early detection of mental health conditions that might otherwise complicate chronic disease management.

Technological innovations, particularly telehealth, represent a promising avenue for expanding integration reach. Amerio et al. (2020) and Hoffmann et al. (2019) note that while digital tools can reduce geographic and stigma-related barriers, they require parallel investment in training and infrastructure to be effective.

Despite the compelling evidence base, the literature reveals persistent structural and systemic barriers. These include misaligned policy priorities, stigma surrounding mental illness, and siloed funding streams (Fusar-Poli et al., 2020; Fox et al., 2012). Overcoming these requires coordinated action between healthcare providers, policymakers, and community stakeholders.

Future research should focus on cost-effectiveness analyses, long-term patient outcomes, and the scalability of integration models in diverse settings. Kantharia et al. (2025) emphasize the importance of synthesizing global lessons to inform contextually appropriate implementation strategies. Importantly, integration efforts must remain patient-centered, ensuring that mental health services are accessible, acceptable, and responsive to the populations they serve.

In conclusion, integrating mental health into family medicine offers a promising path to holistic, equitable care. While challenges remain in financing, training, infrastructure, and cultural adaptation, the cumulative evidence indicates that integrated models can yield substantial benefits for patients, providers, and health systems alike. Sustained commitment to implementation science, workforce development, and policy alignment will be critical for realizing the full potential of this transformative approach.

Conclusion

The integration of mental health services into family medicine practices is a strategic and necessary evolution in primary care delivery. Evidence from diverse contexts demonstrates that integrated models improve patient outcomes, enhance provider collaboration, and contribute to more holistic, person-centered care. This approach not only addresses the gap between physical and mental health services but also offers opportunities for early intervention, continuity of care, and improved management of comorbid conditions. When effectively implemented, integration can mitigate systemic inefficiencies, reduce stigma, and foster equity in health access.

Despite these promising findings, successful implementation requires overcoming entrenched barriers such as funding constraints, workforce shortages, and limited provider training in Behavioral health. Integration must be supported by sustainable financing models, robust interprofessional collaboration, culturally tailored approaches, and scalable innovations like telehealth. Policy commitment, alongside continued research and quality measurement, will be essential to fully realize the benefits of integrated primary and mental healthcare.

Limitations

This review is limited by the heterogeneity of the included studies in terms of populations, integration models, and outcome measures, which precluded meta-analysis and limited direct comparability. Additionally, while the search included diverse geographic settings, there remains an overrepresentation of high-income countries, potentially limiting the generalizability of findings to low-resource contexts. The exclusion of non-English publications may have resulted in language bias. Furthermore, as the review synthesized only published peer-reviewed literature, there is a possibility of publication bias favouring studies with positive outcomes. Finally, variations in the operational definitions of "integration" across studies may affect the interpretation and synthesis of results.

References

1. Aggarwal, M., Knifed, E., Howell, N. A., & Papneja, P. (2020). A qualitative study on the barriers to learning in a primary care-behavioral health integration program in an academic hospital: The family medicine perspective. *Academic Psychiatry*, 44(1), 55-63.
2. Amerio, A., Bianchi, D., Santi, F., Costantini, L., Odone, A., Signorelli, C., et al. (2020). COVID-19 pandemic impact on mental health: A web-based cross-sectional survey on a sample of Italian general practitioners. *Acta Biomedica*, 91(2), 83-88.
3. Asarnow, J. R., Rozenman, M., Wiblin, J., & Zeltzer, L. (2015). Integrated medical-behavioral care compared with usual primary care for child and

- adolescent behavioral health. *JAMA Pediatrics*, 169(10), 929-937.
4. Balasubramanian, B. A., Cohen, D. J., & Jetelina, K. K. (2017). Outcomes of integrated behavioral health with primary care. *Journal of the American Board of Family Medicine*, 30(2), 130-139.
5. Basu, S., Berkowitz, S. A., Phillips, R. L., Bitton, A., Landon, B. E., & Phillips, R. S. (2019). Association of primary care physician supply with population mortality in the United States, 2005-2015. *JAMA Internal Medicine*, 179(4), 506-514.
6. Berge, J. M., Trump, L., Trudeau, S., & Utržan, D. S. (2017). Integrated care clinic: Creating enhanced clinical pathways for integrated behavioral health care. *Families, Systems, & Health*, 35(2), 230-240.
7. Breton, M., Maillet, L., Duhoux, A., Malham, S. A., Gaboury, I., Manceau, L. M., et al. (2020). Evaluation of the implementation and associated effects of advanced access in university family medicine groups: A study protocol. *BMC Family Practice*, 21(1), 41-52.
8. Bryan, C. J., & Morrow, C. E. (2009). Impact of behavioral health consultant interventions on patient symptoms and functioning in an integrated family medicine clinic. *Journal of Clinical Psychology*, 65(3), 281-293.
9. Cohen, D. J., Davis, M., & Balasubramanian, B. A. (2015). Integrating behavioral health and primary care: Consulting, coordinating and collaborating among professionals. *Journal of the American Board of Family Medicine*, 28(Suppl 1), S21-S31.
10. Craven, M. A., Cohen, M., & Campbell, D. (1997). Mental health practices of Ontario family physicians. *Canadian Journal of Psychiatry*, 42(9), 943-949.
11. Fox, M. A., Hodgson, J. L., & Lamson, A. L. (2012). Integration: Opportunities and challenges for family therapists in primary care. *Contemporary Family Therapy*, 34(2), 228-240.
12. Fusar-Poli, P., Salazar de Pablo, G., De Micheli, A., Nieman, D. H., Correll, C. U., Kessing, L. V., et al. (2020). What is good mental health? A scoping review. *European Neuropsychopharmacology*, 31, 33-46.
13. Grazier, K. L., & Smiley, M. L. (2016). Overcoming barriers to integrating behavioral health and primary care services. *Journal of Primary Care & Community Health*, 7(4), 242-246.
14. Hager, K. D., Merighi, J., O'Donnell, C., & Remke, S. (2018). Perceptions of interprofessional collaboration with integrated behavioral health providers in a family medicine clinic. *Journal of Interprofessional Education & Practice*, 12, 61-66.
15. Hoffmann, M., Hartmann, M., & Wensing, M. (2019). Opportunities and challenges for integrating mental health specialist video consultations in office-based routine primary care: Cross-sectional qualitative study among family physicians. *Journal of Medical Internet Research*, 21(8), e13382.
16. Kantharia, F. A., Alhejaili, M. A., Alharbi, M. S., Alsharif, Z. F., Alshammari, R. M., & Al-Almutairi, F. S. (2025). Integration of mental health services in family medicine practice: a systematic review. *International Journal of Medicine in Developing Countries*, 9(2), 448-448.
17. Kilbourne, A. M., Irmiter, C., & Capobianco, J. (2008). Improving integrated general medical and mental health services in community-based practices. *Administration and Policy in Mental Health and Mental Health Services Research*, 35(5), 337-345.
18. Kwan, B. M., & Nease, D. E., Jr. (2013). The state of the evidence for integrated behavioral health in primary care. In M. R. Talen & A. B. Valeras (Eds.), *Integrated behavioral health in primary care: Evaluating the evidence, identifying the essentials* (pp. 65-98). Springer.
19. Liang, D., Mays, V. M., & Hwang, W. C. (2018). Integrated mental health services in China: Challenges and planning for the future. *Health Policy and Planning*, 33(1), 107-122.
20. McGorry, P. D., Mei, C., Chanen, A., Hodges, C., Alvarez-Jimenez, M., & Killackey, E. (2022). Designing and scaling up integrated youth mental health care. *World Psychiatry*, 21(1), 61-76.
21. McGough, P. M., Bauer, A. M., & Collins, L. (2016). Integrating behavioral health into primary care. *Population Health Management*, 19(2), 81-87.
22. McGovern, M., Dent, K., & Kessler, R. (2018). A unified model of behavioral health integration in primary care. *Academic Psychiatry*, 42(6), 758-763.
23. Mullin, D. J., Hargreaves, L., & Auxier, A. (2019). Measuring the integration of primary care and behavioral health services. *Health Services Research*, 54(2), 379-389.
24. Noonan, M., Doody, O., Jomeen, J., & O'Regan, A. (2018). Family physicians perceived role in perinatal mental health. *BMC Family Practice*, 19(1), 170.
25. Ramanuj, P., Ferencsik, E., & Docherty, M. (2019). Evolving models of integrated behavioral health and primary care. *Current Psychiatry Reports*, 21(5), 31.
26. Reed, S. J., Shore, K. K., & Tice, J. A. (2016). Effectiveness and value of integrating behavioral health into primary care. *JAMA Internal Medicine*, 176(5), 691-692.
27. Stein, D. J., Benjet, C., Gureje, O., Lund, C., Scott, K. M., Poznyak, V., et al. (2019). Integrating mental health with other non-communicable diseases. *BMJ*, 364, l295.
28. Vogel, M. E., Kanzler, K. E., Aikens, J. E., & Goodie, J. L. (2017). Integration of behavioral health and primary care: Current knowledge and future directions. *Journal of Behavioral Medicine*, 40(1), 69-84.
29. World Health Organization & World Organization of Family Doctors. (2008). *Integrating mental health into primary care: A global perspective*. WHO Press.
30. Zaman, Q. (2024). Integrating mental health service in primary care: Challenges and opportunities. *Medical Journal of Health*, 4(1), 33-41.